Author Index

Agostini, O.
—, Manzini, S. and Fedi, M.
Determination of 1,3-dimethyl-7-isobutylxanthine in guinea pig plasma by high-performance liquid chromatography 288
Araki, T., see Sakai, T. 73
Arcelloni, C., see Fermo, I. 53
Asano, O., see Yamada, K. 243
Azuma, R., see Marunaka, T. 177
Baty, J.D., see Lindsay, R.M. 292
Bellon, G.
— and Borel, J.P.
Quantitative evaluation of collagen fractions separated by acrylamide gel electrophoresis: its application for collagen typing in normal and pathological tissues 81
Benet, L.Z., see Spahn, H. 331
Ben-Hur, E., see Zuk, M.M. 367
Benny, A.G.
— and Brimer, C.M.
Chromatographic procedure for preparation of a high-purity C 1 esterase inhibitor concentrate 363
Björkman, S.
— and Stanski, D.R.
Simultaneous determination of fentanyl and alfentanil in rat tissues by capillary column gas chromatography 95
Blackett, A.
—, Dhillon, S., Cromarty, J.A., Horne, R. and Richards, G.
Rapid and sensitive high-performance liquid chromatographic assay of midazolam and 1-hydroxymidazolam, the major metabolite, in human serum 326
Bolzacchini, E., see Fermo, I. 53
Bonini, P.A., see Fermo, I. 53
Borchardt, R.T., see Scriba, G.K.E. 31
Borel, J.P., see Bellon, G. 81
Brimer, C.M., see Benny, A.G. 363
Brooks, J.B., see Daneshvar, M.I. 248
Browne, T.R., see Mikati, M.A. 305
Bury, R.W., see Fullinfaw, R.O. 131
Canafax, D.M., see Erdman, G.R. 187
Chichila, T.M.P., see Weidolf, L.O.G. 9
Chiu, S.-H.L.
—, Sestokas, E. and Taub, R.
Application of two-dimensional high-performance liquid chromatography in metabolism studies of ivermectin 217
Churchill, F.C., see Williams, S.B. 197
Coen, C.W., see Opacka-Juffry, J. 41
Collette, L.M., see Wenzel, T.J. 149
Cooper, J.K., see Srinivas, N.R. 105
Crane, S.W., see Zuk, M.M. 367
Cromarty, J.A., see Blackett, A. 326
Dahlen, D.T., see Wenzel, T.J. 149
Daneshvar, M.I.
— and Brooks, J.B.
Improved procedure for preparation of pentafluorobenzyl derivatives of carboxylic acids for analysis by gas chromatography with electron-capture detection 248
Davis, B.A.
Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 23
Dhillon, S., see Blackett, A. 326
Drayer, D.E., see Wu, D.-F. 141
Dysken, M.W., see Vatassery, G.T. 312
Edwards, G., see Milton, K.A. 339
Erdmann, G.R.
—, Canafax, D.M. and Giebink, G.S.
High-performance liquid chromatographic analysis of trimethoprim and sulfamethoxazole in microliter volumes of chinchilla middle ear effusion and serum 187
Fedi, M., see Agostini, O. 288
Feibush, B., see Gisch, D.J. 264
Fennessey, P.V., see Scriba, G.K.E. 31
Fermo, I.
—, Rubino, F.M., Bolzacchini, E., Arcelloni, C., Paroni, R. and Bonini, P.A.
Pre-column derivatization of amino acids with N,N-diethyl-2,4-dinitro-5-fluoroaniline and reversed-phase liquid chromatographic separation 53
Fisher, E., see Kruck, T.P.A. 207
Ford, J.M., see Forsyth, D.R. 352
Forsyth, D.R.
—, Ford, J.M., Truman, D.A., Roberts, C.J.C. and Wilcock, G.K.
Determination of tacrine hydrochloride in human serum by chloroform extraction, reversed-phase high-performance liquid chromatography and fluorimetric detection 352
France, N.C.
—, Holland, P.T., McGhie, T.K. and Wallace, M.R.
Measurement of plasma oxalate by capillary

gas chromatography and its validation by isotope dilution mass spectrometry 1
Fujimaki, Y.
—, Sudo, K. and Tachizawa, H.
High-performance liquid chromatographic determination of a new nootropic, N-(2,6-dimethylphenyl)-2-(2-oxo-1-pyrrolidinyl)-acetamide, in human serum and urine 235
Fullinfaw, R.O.
—, Bury, R.W. and Moulds, R.F.W.
Screening procedure for stimulant laxatives in urine using high-performance liquid chromatography with diode array detection 131
Gengo, F.M., see Schuster, D. 318
Giebink, G.S., see Erdman, G.R. 187
Gisch, D.J.
—, Hunter, B.T. and Feibush, B
Shielded hydrophobic phase: a new concept for direct injection analysis of biological fluids by high-performance liquid chromatography 264
González-Martin, G., see Paulos, C. 359
Greenblatt, D.J., see Mikati, M.A. 305
Grgurinovich, N.
— and Matthews, A.
Analysis of erythromycin and roxithromycin in plasma or serum by high-performance liquid chromatography using electrochemical detection 298
Hendrickson, S.M., see Wenzel, T.J. 149
Henion, J.D., see Weidolf, L.O.G. 9
Herzan, L.A., see Vatassery, G.T. 312
Hoffman, N.E., see Rahman, A. 159
Holland, M.L.
—, Uetz, J.A. and Ng, K.T.
Automated capillary gas chromatographic assay using flame ionization detection for the determination of topiramate in plasma 276
Holland, P.T., see France, N.C. 1
Horne, R., see Blackett, A. 326
Hubbard, J.W., see Srinivas, N.R. 105
Hunter, B.R., see Gisch, D.J. 264
Kalow, W., see Kruck, T.P.A. 207
Katayama, K., see Yamada, K. 243
Kohda, Y., see Yamamoto, K. 167
Kojima, C., see Matsubayashi, K. 225
Komatsu, S., see Marunaka, T. 177
Kruck, T.P.A.
—, Teichert-Kuliszewska, K., Fisher, E., Kalow, W. and McLachlan, D.R.
High-performance liquid chromatographic analysis of desferrioxamine. Pharmacokinetic and metabolic studies 207
Kunze, G., see Paulos, C. 359
Kurotori, M., see Marunaka, T. 177
Kushwaha, R.S., see Williams, M.C. 257
Lalka, D., see Schuster, D. 318
LeDuc, B.W., see Mikati, M.A. 305
Lindsay, R.M.
—, McLaren, A.M. and Baty, J.D.
Reversed-phase high-performance liquid chromatographic assay for the determination of the in vitro acetylation of *p*-aminobenzoic acid by human whole blood 292
McGhie, T.K., see France, N.C. 1
McLachlan, D.R., see Kruck, T.P.A. 207
McLaren, A.M., see Lindsay, R.M. 292
Maniwa, M.
—, Umeno, Y. and Marunaka, T.
Determination of a new hypolipidemic agent, 2-(*trans*-4-isobutylcyclohexyl)-2-oxoethyl benzenesulphonate, and its metabolite by high-performance liquid chromatography and gas chromatography–mass fragmentography in human biological materials 282
Manzini, S., Agostini, O. 288
Marko, V.
Determination of stobadine, a novel cardioprotective drug, using capillary gas chromatography with nitrogen–phosphorus detection after its selective solid-phase extraction from serum 269
Marunaka, T., see Maniwa, M. 282
—, Maniwa, M., Matsushima, E., Yoshida, K.-I., Azuma, R., Kurotori, M. and Komatsu, S.
High-performance liguid chromatographic determination of 6-amidino-2-naphthyl [4-(4,5-dihydro-1*H*-imidazol-2-yl) amino]-benzoate dimethanesulphonate and its metabolites in biological fluids 177
Matsubayashi, K.
—, Kojima, C. and Tachizawa, H.
Determination of tranexamic acid in human serum by high-performance liquid chromatography using selective pre-column derivatization with phenyl isothiocyanate 225
Matsushima, E., see Marunaka, T. 177
Matthews, A., see Grgurinovich, N. 298
Midha, K.K., see Srinivas, N.R. 105
Mikati, M.A.
—, Szabo, G.K., Pylilo, R.J., LeDuc, B.W., Browne, T.R. and Greenblatt, D.J.
Improved high-performance liquid chromatographic assay of antipyrine, hydroxy-

methylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine 305
Milton, K.A.
—, Ward, S.A. and Edwards, G.
Determination of halofantrine and its principal metabolite desbutylhalofantrine in biological fluids by reversed-phase high-performance liquid chromatography 339
Mina, M.
— and Rossomando, E.F.
Assay of ATP-sulfurylase activity from rat liver by high-performance liquid chromatography 63
Minawa, M., see Marunaka, T. 177
Modi, M. Woodruff, see Schuster, D. 318
Moulds, R.F.W., see Fullinfaw, R.O. 131
Mutschler, E., see Spahn, H. 331
Nakagawa, F., see Yamamoto, K. 167
Nancollas, G.H., see Singh, R.P. 373
Newman, H.C., see Zuk, M.M. 367
Ng, K.T., see Holland, M.L. 276
Opacka-Juffry, J.
—, Tacconelli, F. and Coen, C.W.
Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using liquid chromatography with electrochemical detection 41
Paredes, J., see Paulos, C. 359
Paroni, R., see Fermo, I. 53
Patchen, L.C., see Williams, S.B. 197
Paulos, C.
—, Paredes, J., Vasquez, I., Kunze, G. and González-Martin, G.
High-performance liquid chromatographic determination of nifurtimox in human serum 359
Pflugmann, G., see Spahn, H. 331
Pylilo, R.J., see Mikati, M.A. 305
Rahman, A.
— and Hoffman, N.E.
High-performance liquid chromatographic determination of fusidic acid in plasma 159
Reh, E.
Determination of digoxin in serum by on-line immunoadsorptive clean-up high-performance liquid chromatographic separation and fluorescence-reaction detection 119
Reidenberg, M.M., see Wu, D.-F. 141
Richards, G., see Blackett, A. 326
Roberts, C.J.C., see Forsyth, D.R. 352
Rosenthal, I., see Zuk, M.M. 367
Rossomando, E.F., see Mina, M. 63
Rubino, F.M., see Fermo, I. 53
Saitoh, Y., see Yamamoto, K. 167
Sakai, T.
—, Takeuchi, Y., Araki, T. and Ushio, K.
Determination of erythrocyte porphyrins by reversed-phase high-performance liquid chromatography using capsule-type silica gels coated with silicone polymer 73
Schuster, D.
—, Woodruff Modi, M., Lalka, D. and Gengo, F.M.
Reversed-phase high-performance liquid chromatographic assay to quantitate diastereomeric derivatives of metoprolol enantiomers in plasma 318
Scriba, G.K.E.
—, Borchardt, R.T., Zirrolli, J.A. and Fennessey, P.V.
Selected-ion monitoring gas chromatographic–mass spectrometric analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 31
Sestokas, E., see Chiu, S.-H.L. 217
Singh, R.P.
— and Nancollas, G.H.
Determination of phosphate, sulfate and oxalate in urine by ion chromatography 373
Spahn, H.
—, Spahn, I., Pflugmann, G. and Mutschler, E.
Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 331
Spahn, I., see Spahn, H. 331
Srinivas, N.R.
—, Hubbard, J.W., Cooper, J.K. and Midha, K.K.
Enantioselective gas chromatographic assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma 105
Stanski, D.R., see Björkman, S. 95
Sudo, K., see Fujimaki, Y. 235
Szabo, G.K., see Mikati, M.A. 305
Tacconelli, F., see Opacka-Juffry, J. 41
Tachizawa, H., see Fujimaki, Y. 235
—, see Matsubayashi, K. 225
Takeuchi, Y., see Sakai, T. 73
Taub, R., see Chiu, S.-H.L. 217
Teichert-Kuliszewska, K., see Kruck, T.P.A. 207

Thongnopnua, P.
— and Zimmerman, C.L.
Simultaneous microassay for etretinate and its active metabolite, etretin, by reversed-phase high-performance liquid chromatography 345
Truman, C.A., see Forsyth, D.R. 352
Tyczkowska, K., see Zuk, M.M. 367
Uetz, J.A., see Holland, M.L. 276
Umeno, Y., see Maniwa, M. 282
Ushio, K., see Sakai, T. 73
Vasquez, I., see Paulos, C. 359
Vatassery, G.T.
—, Herzan, L.A. and Dysken, M.W.
Simultaneous determination of very low concentrations of haloperidol and reduced haloperidol in human serum by a liquid chromatographic method 312
Wallace, M.R., see France, N.C. 1
Ward, S.A., see Milton, K.A. 339
Weidolf, L.O.G.
—, Chichila, T.M.P. and Henion, J.D.
Screening, confirmation and quantification of boldenone sulfate in equine urine after administration of boldenone undecylenate (Equipoise™) 9
Wenzel, T.J.
—, Collette, L.M., Dahlen, D.T., Hendrickson, S.M. and Yarmaloff, L.W.
Liquid chromatographic and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection 149
Wilcock, G.K., see Forsyth, D.R. 352
Williams, M.C.
— and Kushwaha, R.S.
Fractionation of baboon chylomicrons and very-low-density lipoproteins by high-performance liquid chromatography 257
Williams, S.B.
—, Patchen, L.C. and Churchill, F.C.
Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by high-performance liquid chromatography with fluorescence detection 197
Woodruff Modi, M., see Schuster, D. 318
Wu, D.-F.
—, Reidenberg, M.M. and Drayer, D.E.
Determination of gossypol enantiomers in plasma after administration of racemate using high-performance liquid chromatography with precolumn chemical derivatisation 141
Yamada, K.
—, Asano, O., Yoshimura, T. and Katayama, K.
Highly sensitive gas chromatographic–mass spectrometric method for the determination of platelet-activating factor in human blood 243
Yamamoto, K.
—, Kohda, Y., Saitoh, Y. and Nakagawa, F.
Determination of ambenonium in biological samples by reversed-phase ion-pair liquid chromatography 167
Yarmaloff, L.W., see Wenzel, T.J. 149
Yoshida, K.-I., see Marunaka, T. 177
Yoshimura, T., see Yamada, K. 243
Zimmerman, C.L., see Thongnopnua, P. 345
Zirrolli, J.A., see Scriba, G.K.E. 31
Zuk, M.M.
—, Tyczkowska, K., Ben-Hur, E., Newman, H.C., Rosenthal, I. and Crane, S.W.
Reversed-phase liquid chromatographic determination of chloroaluminum phthalocyanine tetrasulfonate in canine tissues and fluids 367

Subject Index

Alfentanil
Simultaneous determination of fentanyl and alfentanil in rat tissues by capillary column GC 95

Ambenonium
Determination of ambenonium in biological samples by reversed-phase ion-pair LC 167

6-Amidino-2-naphthyl [4-(4,5-dihydro-1*H*-imidazol-2-yl)amino]benzoate dimethanesulphonate
HPLC determination of 6-amidino-2-naphthyl [4-(4,5-dihydro-1*H*-imidazol-2-yl)amino]benzoate dimethanesulphonate and its metabolites in biological fluids 177

Amines
Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 23

Amino acids
Pre-column derivatization of amino acids with N,N-diethyl-2,4-dinitro-5-fluoroaniline and reversed-phase LC separation 53

p-Aminobenzoic acid
Reversed-phase HPLC assay for the determination of the in vitro acetylation of *p*-aminobenzoic acid by human whole blood 292

Antipyrine
Improved HPLC assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine 305

ATP-sulfurylase
Assay of ATP-sulfurylase activity from rat liver by HPLC 63

Bisacodyl
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131

Bisacodyl diphenol
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131

Bisdesethylchloroquine
Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 197

Bisoxatin
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131

Bisoxatin diphenol
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131

Boldenone sulfate
Screening, confirmation and quantification of boldenone sulfate in equine urine after administration of boldenone undecylenate (Equipoise™) 9

Carboxylic acids
Improved procedure for preparation of pentafluorobenzyl derivatives of carboxylic acids for analysis by GC with electron-capture detection 248

Carprofen
Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 331

Catecholamines
Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 23

Catecholamines
Selected-ion monitoring GC–MS analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 31

Catecholamines
Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 41

Chloroaluminum phthalocyanine tetrasulfonate
Reversed-phase LC determination of chloroaluminum phthalocyanine tetrasulfonate in canine tissues and fluids 367

Chylomicrons
Fractionation of baboon chylomicrons and very-low-density lipoproteins by HPLC 257

Collagen
Quantitative evaluation of collagen fractions separated by acrylamide gel electrophoresis: its application for collagen typing in normal and pathological tissues 81

Danthron
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131
Desbutylhalofantrine
Determination of halofantrine and its principal metabolite desbutylhalofantrine in biological fluids by reversed-phase HPLC 339
Desethylchloroquine
Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 197
Desethylhydroxychloroquine
Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 197
Desferrioxamine
HPLC analysis of desferrioxamine. Pharmacokinetic and metabolic studies 207
N,N-Diethyl-2,4-dinitro-5-fluoroaniline
Pre-column derivatization of amino acids with N,N-diethyl-2,4-dinitro-5-fluoroaniline and reversed-phase LC separation 53
Digoxin
Determination of digoxin in serum by on-line immunoadsorptive clean-up HPLC separation and fluorescence-reaction detection 119
3,4-Dihydroxybenzylamine
Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 41
1,3-Dimethyl-7-isobutylxanthine
Determination of 1,3-dimethyl-7-isobutylxanthine in quinea pig plasma by HPLC 288
N-(2,6-Dimethylphenyl)-2-(2-oxo-1-pyrrolidinyl)acetamide
HPLC determination of a new nootropic, N-(2,6-dimethylphenyl)-2-(2-oxo-1-pyrrolidinyl)acetamide, in human serum and urine 235
Dopamine
Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 41
Enzymes
Assay of ATP-sulfurylase activity from rat liver by HPLC 63
Epinephrine
Selected-ion monitoring GC–MS analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 31
Epinephrine
Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 41
Epinine
Selected-ion monitoring GC–MS analysis of catecholamines: enhancement of sensitivity by a simple clean-up step on Sephadex G-10 31
Erythromycin
Analysis of erythromycin and roxithromycin in plasma or serum by HPLC using electrochemical detection 298
C 1 Esterase inhibitor
Chromatographic procedure for preparation of a high-purity C 1 esterase inhibitor concentrate 363
Etretin
Simultaneous microassay for etretinate and its active metabolite, etretin, by reversed-phase HPLC 345
Etretinate
Simultaneous microassay for etretinate and its active metabolite, etretin, by reversed-phase HPLC 345
Europium(III)
LC and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection 149
dl-Fenfluramine
Enantioselective GC assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma 105
Fentanyl
Simultaneous determination of fentanyl and alfentanil in rat tissues by capillary column GC 95
Fusidic acid
HPLC determination of fusidic acid in plasma 159
Gossypol
Determination of gossypol enantiomers in plasma after administration of racemate using HPLC with precolumn chemical derivatisation 141

Halofantrine
Determination of halofantrine and its principal metabolite desbutylhalofantrine in biological fluids by reversed-phase HPLC 339
Haloperidol
Simultaneous determination of very low concentrations of haloperidol and reduced haloperidol in human serum by a LC method 312
4-Hydroxyantipyrine
Improved HPLC assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine 305
Hydroxychloroquine
Analysis of blood and urine samples for hydroxychloroquine and three major metabolites by HPLC with fluorescence detection 197
Hydroxymethylantipyrine
Improved HPLC assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine 305
1-Hydroxymidazolam
Rapid and sensitive HPLC assay for midazolam and 1-hydroxymidazolam, the major metabolite, in human serum 326
Indoleamines
Effects of long-term storage on the concentrations of the unconjugated acidic metabolites of the trace amines, indoleamines and catecholamines 23
2-(*trans*-4-Isobutylcyclohexyl)-2-oxoethyl benzensulphonate
Determination of a new hypolipidemic agent, 2-(*trans*-4-isobutylcyclohexyl)-2-oxoethyl benzenesulphonate, and its metabolite by HPLC and GS–mass fragmentography in human biological materials 282
Ivermectin
Application of two-dimensional HPLC in metabolism studies of ivermectin 217
Laxatives
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131
L-Leucinamide
Measurement of carprofen enantiomer concentrations in plasma and urine using L-leucinamide as the chiral coupling component 331
Lipoproteins
Fractionation of baboon chylomicrons and very-low-density lipoproteins by HPLC 257
Metoprolol
Reversed-phase HPLC assay to quantitate diastereomeric derivatives of metoprolol enantiomers in plasma 318
Midazolam
Rapid and sensitive HPLC assay for midazolam and 1-hydroxymidazolam, the major metabolite, in human serum 326
Nifurtimox
HPLC determination of nifurtimox in human serum 359
Norantipyrine
Improved HPLC assay of antipyrine, hydroxymethylantipyrine, 4-hydroxyantipyrine and norantipyrine in urine 305
Norepinephrine
Sensitive method for determination of picogram amounts of epinephrine and other catecholamines in microdissected samples of rat brain using LC with electrochemical detection 41
dl-Norfenfluramine
Enantioselective GC assay with electron-capture detection for *dl*-fenfluramine and *dl*-norfenfluramine in plasma 105
Oxalate
Measurement of plasma oxalate by capillary GC and its validation by isotope dilution MS 1
Oxalate
Determination of phosphate, sulfate and oxalate in urine by ion chromatography 373
Phenolphthalein
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131
Phosphate
Determination of phosphate, sulfate and oxalate in urine by ion chromatography 373
Phospholipids
Highly sensitive GC–MS method for the determination of platelet-activating factor in human blood 243
Platelet-activating factor
Highly sensitive GC–MS method for the determination of platelet-activating factor in human blood 243
Porphyrins
Determination of erythrocyte porphyrins by reversed-phase HPLC using capsule-type

silica gels coated with silicone polymer 73
Protoporphyrins
Determination of erythrocyte porphyrins by reversed-phase HPLC using capsule-type silica gels coated with silicone polymer 73
Reduced haloperidol
Simultaneous determination of very low concentrations of haloperidol and reduced haloperidol in human serum by a LC method 312
Rhein
Screening procedure for stimulant laxatives in urine using HPLC with diode array detection 131
Roxithromycin
Analysis of erythromycin and roxithromycin in plasma or serum by HPLC using electrochemical detection 298
Shielded hydrophobic phase
Shielded hydrophobic phase: a new concept for direct injection analysis of biological fluids by HPLC 264
Stobadine
Determination of stobadine, a novel cardioprotective drug, using capillary GC with nitrogen–phosphorus detection after its selective solid-phase extraction from serum 269
Sulfamethoxazole
HPLC analysis of trimethoprim and sulfamethoxazole in microliter volumes of chinchilla middle ear effusion and serum 187
Sulphate
Determination of phosphate, sulfate and oxalate in urine by ion chromatography 373
Tacrine hydrochloride
Determination of tacrine hydrochloride in human serum by chloroform extraction, reversed-phase HPLC and fluorimetric detection 352
Tetracycline
LC and flow injection analysis of tetracycline using sensitized europium(III) luminescence detection 149
Topiramate
Automated capillary GC assay using flame ionization detection for the determination of topiramate in plasma 276
Tranexamic acid
Determination of tranexamic acid in human serum by HPLC using selective precolumn derivatization with phenyl isothiocyanate 225
Trimethoprim
HPLC analysis of trimethoprim and sulfamethoxazole in microliter volumes of chinchilla middle ear effusion and serum 187
Very-low-density lipoproteins
Fractionation of baboon chylomicrons and very-low-density lipoproteins by HPLC 257